

Problem 1


(a)

(i) $1 - i$


$$z = 1 - i \quad |z| = \sqrt{1^2 + (-1)^2} \\ = \sqrt{2}$$

$$\begin{aligned}\operatorname{Arg}(z) &= \tan^{-1} \frac{-1}{1} \\ &= 2\pi - \frac{\pi}{4} \\ &= \frac{7\pi}{4}\end{aligned}$$

$$z = \sqrt{2} e^{\frac{7\pi}{4}i}$$



$$\begin{aligned}(\text{ii}) \quad z &= 2 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) \quad z = 2e^{\frac{\pi}{6}i} \\ &= \frac{2\sqrt{3}}{2} + i \frac{2}{2} \\ &= \sqrt{3} + i\end{aligned}$$



$$(\text{iii}) \quad z = \sqrt{-a - bi}$$


$$(iv) \quad z = \sqrt[6]{1}, \quad 0 < \theta < \frac{\pi}{2}$$

$$z^6 = 1$$

$$z = e^{\frac{2\pi k}{6}i}, \quad k = 0, 1, \dots, 5$$

$$\Rightarrow z = e^{\frac{\pi}{3}i} \\ = \frac{1 + \sqrt{3}i}{2}$$

$$z = \frac{1}{2} + \frac{\sqrt{3}}{2}i \quad z = e^{\frac{\pi}{3}i}$$



$$(v) \quad z = \left(\frac{1+i}{\sqrt{2}} \right)^{-13}$$


$$= \left(\frac{\sqrt{2}}{1+i} \right)^{13}$$

$$= \frac{-\sqrt{2} + \sqrt{2}i}{2}$$

$$\frac{\sqrt{2}}{1+i} = \frac{\sqrt{2} - \sqrt{2}i}{2} \quad \theta = \tan^{-1} \frac{-1}{1} \\ = e^{\frac{7\pi}{4}i} \quad |z| = 1$$

$$\left(e^{\frac{7\pi}{4}i} \right)^{13} = e^{\frac{91\pi}{4}i}$$

$$= e^{\frac{3\pi}{4}i}$$



$$(b) z^4 + 4 = 0$$

$$z^2 + 2z + 2 = 0$$

$$\text{Let } x = z^2,$$

$$\text{then } x^2 + 4 = 0$$

$$x = \frac{\pm \sqrt{-16}}{2}$$

$$= \pm 2i$$

$$z = \frac{-2 \pm \sqrt{4 - 8}}{2}$$

$$= -1 \pm i$$

$$= \sqrt{2} e^{\frac{3\pi}{4}i}, \sqrt{2} e^{-\frac{3\pi}{4}i}$$

$$\Rightarrow z^2 = 2i, -2i \\ = 2e^{\frac{\pi}{2}i}, 2e^{-\frac{\pi}{2}i}$$

$$z = \sqrt{2} e^{\frac{\pi}{4}i}, \sqrt{2} e^{-\frac{\pi}{4}i}$$

$$z = e^{\frac{\frac{\pi}{2}+2k\pi}{2}i} \text{ where } k=0,1 \\ = e^{(\frac{\pi}{4}+k\pi)i} \quad w = e^{ki}$$

$$\Rightarrow z = \sqrt{2} e^{\frac{\pi}{4}i}, \sqrt{2} e^{\frac{5\pi}{4}i}, \sqrt{2} e^{-\frac{3\pi}{4}i}, \sqrt{2} e^{\frac{7\pi}{4}i}$$